



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

**Pseudatrichia pilosa** Coq.

The hairs of the thorax and femora are unusually long, a very marked character, especially with fresh, well preserved specimens.

**Pseudatrichia flaviceps** Coq.

A well marked species, easily distinguished by the two large yellow spots on the anterior margin of the thorax, and the entire yellow front and face. The whitish line below each wing is very narrow, sometimes hardly visible.

In looking over this species in the National Museum I discovered a male, and as this sex has not been described, Dr. Coquillett kindly gave me permission to do so.

Front very narrow, as wide as the second antennal joint at its narrowest part; antennæ yellow, but the third joint brownish; the yellow of the thorax less extended, limited to lateral humeral angles; the postalar calluses entirely yellow (those of the female brownish); the knobs of halteres entirely white. This specimen the color in general is more brownish, and the markings more whitish; the legs accordingly, so that the yellow of the knees is not so marked. Length 3 mm.

One specimen from Williams, Arizona (Barber). Collection of U. S. N. M.

**Pseudatrichia longurio** Lw.

*Atrichia* Loew, Cent., vii, 76, 1866.

"Male.—Black, opaque; third antennal joint dark rufous, the anterior edge darker; humeri and scutellum yellowish or brownish; feet and coxæ brownish; but the extreme bases of coxæ and apex of tibiæ and tarsi darker brown; abdomen slender, elongated; hypopygium dark rufous; wings grayish-hyaline, veins brownish. Length 4 lin."

*Hab.*—Mexico.

## EXPLANATION OF PLATE II.

- Fig. 1. *Scenopinus fenestralis*, wing.  
 " 2. " *nubilipes*, ♂.  
 " 3. *Metatrichia bulbosus*, ♂.  
 " 4. " " wing.  
 " 5. *Pseudatrichia griseola*, ♀.  
 " 6. " *unicolor*, ♂.  
 " 7. " " wing.  
 " 8. " " (variety), ♀.  
 " 9. " " wing.

## CORRECTION TO PAPER I.—ORTALIDÆ.

The table on page 285 (Trans., Vol. 32), at 2a, should read:

- 2a. Antennæ at least six times as long as the width of the second joint; third joint elongated, unusually long ..... **Stictomyia** Big.  
 Antennæ not so long; third joint more or less oval ..... 3 of manual.